

VERSION WITH MARKINGS TO SHOW CHANGES

IN THE SPECIFICATION:

Page 5, amend the paragraph beginning on line 10 to read as follows:

- performing on the same sample a second thermogram in the same range and under a pressure Ph of a hydrocarbon gas, T1 being a temperature low enough to obtain the formation of hydrates in the sample at a gas pressure Ph, P2-T2 being high enough to obtain hydrate dissociation,

IN THE CLAIMS:

1. (Amended) A method for determining gas hydrate formation conditions in a well fluid, comprising the following stages :

- taking a fluid sample,
- placing this sample in a calorimetry cell,
- performing on this sample a reference thermogram in a temperature range between T1 and T2,
- performing on the same sample a second thermogram in the same range and under a pressure Ph of a hydrocarbon gas, T1 being a temperature low enough to obtain the formation of hydrocarbon gas hydrates in the sample at a gas pressure Ph, P2-T2 being high enough to obtain hydrate dissociation,
- identifying a peak in the second thermogram corresponding to the hydrocarbon gas hydrates dissociation zone and deducing therefrom a hydrocarbon gas hydrates

dissociation temperature,

- determining the hydrocarbon gas hydrate formation conditions for the fluid considered.

2. (Amended) A method as claimed in claim 1, wherein pressure P_h is determined as a function of the pressure of the well fluid close to the zones where the appearance of hydrocarbon gas hydrates is critical.

IN THE ABSTRACT:

Method of determining the gas hydrate formation conditions in a well fluid, comprising the following stages :

- taking a fluid sample,
- placing this sample in a calorimetry cell,
- performing on this sample a reference thermogram in a temperature range between T_1 and T_2 ,
- performing on the same sample a second thermogram in the same range and under a pressure P_h of a hydrocarbon gas, T_1 being a temperature low enough to obtain the formation of hydrates in the sample at a gas pressure P_h , P_2 - T_2 being high enough to obtain hydrate dissociation,

identifying a peak in the second thermogram corresponding to the hydrates dissociation zone and deducing therefrom a hydrates dissociation temperature.